

Weekly COVID-19 Vaccine Updates

Number 13, 10 June 2021



Introduction

This document summarises the vaccine efficacy and effectiveness, the vaccine specifications, the vaccine development pipeline and the timeline for World Health Organization (WHO) review of the various COVID-19 vaccines in late phase development. This document is updated weekly.

- Vaccine efficacy refers to the performance of a vaccine in a controlled clinical trial (study) situation
- Vaccine effectiveness refers to the performance of a vaccine in a population under real-world conditions

Key messages









- COVID-19 vaccine efficacy results from different trials cannot be directly compared against each other. They must be interpreted in the context of study designs (including case definitions, clinical endpoints, access to testing), target populations, and COVID-19 epidemiologic conditions (including circulation of variants of concern)
- All COVID-19 vaccines in late phase development report high vaccine efficacy against severe COVID-19 and favourable safety profiles
- Pfizer/BioNTech and AstraZeneca both show high vaccine effectiveness in the UK and Israel where the B.1.1.7 (UK) variant is circulating. Both vaccines have shown they are similarly effective against transmission in UK. Sinovac has shown high vaccine effectiveness in Chile where the P1 and B.1.1.7 variants are circulating. Sinopharm has shown high vaccine effectiveness in Bahrain.
- The US FDA, UK MHRA, EU EMA and Health Canada have authorised the Pfizer/BioNTech vaccine for emergency use in adolescents aged 12-15 years¹⁻⁴
- Mixed vaccine schedules (i.e. delivering different types of vaccine for the first and second dose) are under investigation as these could facilitate better protection against variants of concern and enable vaccination programs to continue if a particular vaccine is unavailable.
- A very rare clotting disorder with low platelets (Thrombosis with Thrombocytopenia Syndrome – TTS) has been associated with the AstraZeneca and Johnson & Johnson vaccines.⁵⁻⁷ The majority of cases fully recover with adequate treatment. The risk following the first dose of AstraZeneca vaccine has been estimated by the EMA as 1 in 100,000 and by the Australian Technical Advisory Group on Immunisation (ATAGI) as 1 in 54,000.^{8,9} Risk of TTS is much lower following the *second* dose of AstraZeneca vaccine: estimate in the UK is 1 in 769,000 second doses.¹⁰
- The risk of TTS following the first dose of Johnson & Johnson vaccine has been estimated as 1 in 319,000 in the USA¹¹
- Appropriate communication on the benefit-risk profile of COVID-19 vaccines (Page 12) remains crucial to maintain confidence in immunisation programmes and to avoid vaccine hesitancy

New updates

Key updates include (*also highlighted in yellow text in the document*):

- Vaccine effectiveness of Pfizer/BioNTech in Denmark (Pages 5 and 7):
 - All priority groups:
 - Infection: 82% (79-84)
 - Hospitalisation: 93% (89-96)
 - Death: 94% (90-96)
 - ≥85 years:
 - Infection: 77% (50-89)
 - Comorbidities with high risk of severe COVID-19:
 - Infection: 71% (58-80)
 - Hospitalisation: 81% (49-93)
- Vaccine effectiveness against B.1.617.2 (Delta variant) in the UK (updated with data following a single dose):
 - AstraZeneca:
 - ≥21 days after first dose: 32.9% (19.3-44.3)
 - ≥14 days after second dose: 59.8% (28.9-77.3)
 - Pfizer/BioNTech:
 - ≥21 days after first dose: 33.2% (8.3-51.4)
 - ≥14 days after second dose: 87.9% (78.2-93.2)
- Vaccine effectiveness against B.1.1.7 (Alpha variant) in the UK:
 - AstraZeneca:
 - ≥21 days after first dose: 51.4% (47.3 to 55.2)
 - ≥14 days after second dose: 66.1% (54.0-75.0)
 - Pfizer/BioNTech:
 - ≥21 days after first dose: 49.2% (42.6 to 55.0)
 - ≥14 days after second dose: 93.4% (90.4-95.5)
- A new table has been added on mixed dose vaccine safety and immune responses (Page 9)
- The Pfizer/BioNTech vaccine has been approved for use in 12-15 year olds in the UK
- A case series has been published on myocarditis/myopericarditis within 4 days of the second dose of Pfizer/BioNTech vaccine in 7 males aged 14-19 years in the USA. The risk of myocarditis has been estimated between 1 in 3,000 to 1 in 6,000 following the second dose of Pfizer/BioNTech vaccine in men aged 16-24 years in Israel. Cases were mostly mild and resolved with treatment (Page 10)
- The adverse events following immunisation table has been updated with Sinovac data (Page 10)
- Potential risk of TTS following AstraZeneca and Johnson & Johnson vaccines in PICs has been updated with latest data from Australia and USA (Page 12)

COVID-19 Vaccine Specifications

	ASTRAZENECA	GAMALEYA	JOHNSON & JOHNSON	MODERNA	NOVAVAX	PFIZER/BIONTECH	SINOVAC	SINOPHARM
VACCINE TYPE	Viral vector (chimpanzee adenovirus ChAdOx1)	Viral vector (recombinant adenovirus types 5 and 26)	Viral vector (recombinant adenovirus type 26)	mRNA	Protein subunit	mRNA	Inactivated virus	Inactivated virus
Available Through COVAX	✓	-	✓	-	✓	✓	-	-
Doses Required	 8-12 weeks apart* 4 weeks apart (Product Information)	 3 weeks apart		 4 weeks apart*	 3 weeks apart	 3 weeks apart*	 2 weeks apart	 3 weeks apart
Shipping, Storage & Presentation	Normal cold chain requirements (2-8°C); 10-dose vials	-18.5°C (liquid form); 2-8°C (dry form)	Shipped at -20°C; 2-8°C for up to 3 months; 5-dose vials	-25°C to -15°C; 10-dose vials	2-8°C; 10-dose vials	-80°C to -60°C; 2-8°C for up to 1 month; 6-dose vials	2-8°C; Single-dose vials	2-8°C; Single-dose vials/pre-filled syringes
Approval by a Stringent Regulatory Authority (SRA)	WHO EUL, EMA, TGA, MHRA	Under review by WHO SAGE	WHO EUL, EMA, FDA, MHRA	WHO EUL, EMA, FDA	Under review by WHO SAGE	WHO EUL, EMA, FDA, TGA, MHRA	WHO EUL	WHO EUL

*Based on WHO Strategic Advisory Group of Experts on Immunization (SAGE) recommendations

WHO EUL: WHO Emergency Use Listing
EMA: European Medicines Agency
FDA: Food and Drug Administration (US)
TGA: Therapeutic Goods Administration (Australia)
MHRA: Medicines and Healthcare Products Regulatory Agency (UK)

COVID-19 Vaccine Efficacy

VACCINE	VACCINE EFFICACY			
	SYMPTOMATIC INFECTION	MODERATE-SEVERE	SEVERE	HOSPITALISATION/DEATH
AstraZeneca	UK: 66.7% (57.4-74.0) ¹² USA, Chile, Peru: 76% ¹³ (not peer-reviewed) Single dose in UK (22-90 days post-vaccination): 76.0% (59.3 to 85.9) ¹² Efficacy with different interval between doses in UK: 12+ weeks: 82.4% (2.7-91.7) <6 weeks: 54.9% (32.7-69.7) ¹²	-	Severe/critical and hospitalisation in USA, Chile, Peru: 100% ¹³ (not peer-reviewed) UK: 100% (15 cases in the placebo group) ¹²	Hospitalisation in UK: 100% (9 cases in placebo group) ¹²
Gamaleya	91.6% (85.6-95.2) ¹⁴	Moderate-severe: 100% (20 cases in the placebo group) ¹⁴	-	-
Johnson & Johnson	-	Moderate to severe/critical: All sites: 66.1% (55.0-74.8) USA: 72.0% (58.2-81.7) Latin America: 61.0% (46.9-71.8) South Africa: 64.0% (41.2-78.7) ¹⁵	85.4% (54.2-96.9) ¹⁵	100% (7 deaths in placebo group) ¹⁵
Moderna	USA: 94.1% (89.3-96.8) ¹⁶ USA: >90% ¹⁷ 12-17 years in USA: 93% (1 case in vaccine arm) ¹⁸	-	USA: 100% (30 cases in placebo group) ¹⁶ US: >95% ¹⁷	USA: 100% (1 death in placebo group) ¹⁶
Novavax	UK: 89.7% (80.2-94.6) ¹⁹	-	-	-
Pfizer/BioNTech	Argentina, Brazil, Germany, South Africa, Turkey and the USA: 94.6% (89.9-97.3) ²⁰ 12-15 years in USA: 100% ²¹	-	Argentina, Brazil, Germany, South Africa, Turkey and the USA: 88.9% (20.1-99.7) ²⁰	-
Sinovac	Brazil: 50.7% (35.9-62.0) Chile: 67% (65-69) Indonesia: 65% (20-85) Turkey: 84% (65-92) ²²	Requiring medical assistance in Brazil: 83.7% (58.0-93.7) Moderate-severe: 100% (56.4-100.0) ²³	Brazil: Moderate-severe: 100% (56.4-100.0) ²³	Hospitalisation: Brazil: 100% (56-100) Chile: 85% (83-97) Turkey: 100% (20-100) ²²
Sinopharm	UAE, Bahrain, Egypt and Jordan: 78.1% (64.9-86.3) ²²	-	-	Hospitalisation in UAE, Bahrain, Egypt and Jordan: 78.7% (26.0-93.9) ²²

COVID-19 Vaccine Effectiveness

VACCINE	SEVERE	HOSPITALISATION / DEATH	OTHER OUTCOMES
AstraZeneca	-	<p>Single dose in Scotland: 94% (73-99)²⁴</p> <p>Risk of death in vaccine failures compared to unvaccinated cases in England reduced by: 55% (41-66)²⁵ (not peer reviewed)</p> <p>Single dose against hospitalisation in Spain: 92% (46-99)²⁶</p>	<p>Pooled analysis of Pfizer/BioNTech and AstraZeneca vaccines in elderly care home residents in UK: Reduction in risk of infection 4 weeks after-single dose: 56% Reduction in risk of infection 5 weeks after single dose: 62%²⁷</p> <p>Pooled analysis of Pfizer/BioNTech and AstraZeneca vaccines: reduced odds of infection post-second dose: 70% (62-77)²⁸</p> <p>Single dose in Spain: Any infection: 44% (31-54); Symptomatic infection: 50% (37-61)²⁶</p>
Moderna	-	-	<p>Pooled analysis of Moderna and Pfizer/BioNTech vaccines in USA: Infections in nonvaccinated: 234 of 8969; 2.61% (2.29-2.96) Fully vaccinated: 4/8121; 0.05% (0.01-0.13)²⁹</p> <p>Pooled analysis of Moderna and Pfizer/BioNTech vaccines in USA: Fully vaccinated: 90% (68-97) Two weeks after first dose: 80% (59-90)³⁰</p>
Pfizer/BioNTech	<p>Israel: 92% (75-100)³¹</p> <p>Severe/critical in Israel: 97.5% (97.1-97.8)³²</p>	<p>Single dose in Scotland: 85% (76-91)²⁴</p> <p>Risk of death in vaccine failures compared to unvaccinated cases in England reduced by: Single dose: 44% (32-53)</p> <p>Fully vaccinated: 69% (31-86)²⁵ (not peer reviewed)</p> <p>Israel: Hospitalisation: 97.2% (96.8-97.5); Death: 96.7% (96.0-97.3)³²</p> <p>Hospitalisation in Spain: 94% (60-99)²⁶</p> <p>Priority groups in Denmark: Hospitalisation: 93% (89-96); Death: 94% (90-96)³³</p>	<p>Pooled analysis of Moderna and Pfizer/BioNTech vaccines in USA: Infections in nonvaccinated: 234 of 8969; 2.61% (2.29-2.96) Fully vaccinated: 4/8121; 0.05% (0.01-0.13)²⁹</p> <p>Pooled analysis of Moderna and Pfizer/BioNTech vaccines in USA: Fully vaccinated: 90% (68-97) Two weeks after first dose: 80% (59-90)³⁰</p> <p>Symptomatic infection in Israel: 94% (87-98)³¹ Any infection in Israel: 90% (79-95)³⁴</p> <p>Israel: Any infection: 95.3% (94.9-95.7); Symptomatic infection: 97.0% (96.7-97.2)³²</p> <p>Pooled analysis of Pfizer/BioNTech and AstraZeneca vaccines in elderly care home residents in UK: Reduction in risk of infection 4 weeks after-single dose: 56% Reduction in risk of infection 5 weeks after single dose: 62%²⁷</p> <p>Documented infection in Israel: incidence decreased from 9.4 infections per 1,000 HCWs in the week following first dose to <1.0 infection per 1,000 HCWs per week from 1 week after the second dose³⁵</p> <p>Pooled analysis of Pfizer/BioNTech and AstraZeneca vaccines: reduced odds of infection post-second dose: 70% (62-77)²⁸</p> <p>Spain: Any infection: 65% (56-73); Symptomatic infection: 82% (73-88)²⁶</p> <p>Infection in priority groups in Denmark: 82% (79-84)³³</p>
Sinovac	-	<p>Chile: Hospital admission: 85% (83-87); ICU admission: 89% (84-92); Death: 80% (73-86)³⁶ (not peer reviewed)</p>	<p>Symptomatic infection in Chile: 67% (65-69)³⁶ (not peer reviewed)</p>
Sinopharm	-	-	<p>Symptomatic infection in Bahrain: 90% (88-91)²²</p>

Vaccine Efficacy/Effectiveness Against Variants

Refer to previous table for vaccine effectiveness results for the Pfizer/BioNTech vaccine in Scotland, England and Israel, where all locations had predominant B.1.1.7 circulation. There are four Variants of Concern listed by WHO.³⁷ The WHO recommends labelling SARS-CoV-2 variants with letters of the Greek alphabet, as in the table below.³⁸

VACCINE	VACCINE EFFICACY/EFFECTIVENESS							
	B.1.1.7 (ALPHA) VARIANT		B.1.351 (BETA) VARIANT		P.1 (GAMMA) VARIANT		B.1.617.2 (DELTA) VARIANT	
	ANY INFECTION	SEVERE	ANY INFECTION	SEVERE	ANY INFECTION	SEVERE	ANY INFECTION	SEVERE
AstraZeneca	70.4% (43.6-84.5) (vs. 81.5% (67.9-89.4) against wild variant in UK) ³⁹ Effectiveness: ≥21 days after one dose: 51.4% (47.3 to 55.2); ≥14 days after two doses: 66.1% (54.0-75.0) ⁴⁰	-	10.4% (-76.8 to 54.8) ⁴¹	Study underway ¹⁵	-	-	Effectiveness: ≥21 days after one dose: 32.9% (19.3-44.3); ≥14 days after second dose: 59.8% (28.9-77.3) ⁴⁰	
Johnson & Johnson	-	-	-	Moderate to severe/critical: 64.0% (41.2-78.7) Severe/critical: 81.7% (46.2-95.4) ¹⁵	-	Moderate to severe/critical: 68.1% (48.8-80.7) Severe/critical: 87.6% (7.8-99.7) ¹⁵		
Novavax	86.3% (71.3-93.5) (vs. 96.4% (73.8-99.5) against wild variant in UK) ¹⁹	-	South Africa: 51.0% (-0.6 to 76.2) ⁴²	-	-	-		
Pfizer/BioNTech	Case-control study in Israel: Vaccinees infected between 2 weeks after the first dose and 1 week after the second dose, were disproportionately infected with B.1.1.7 (odds ratio of 26:10) ⁴³ Effectiveness in Qatar: 89.5% (85.9-92.3) ⁴⁴ Effectiveness: ≥21 days after one dose: 49.2% (42.6 to 55.0) ≥14 days after second dose: 93.4% (90.4-95.5) ⁴⁰	Effectiveness in Qatar: 100% (81.7-100) ⁴⁴	Israel case-control study: Vaccinees infected at least 1 week after the second dose were disproportionately infected with B.1.351 (odds ratio: 8:1) ⁴³ Effectiveness in Qatar: 75.0% (70.5-78.9) ⁴⁴	Effectiveness in Qatar: 100% (73.7-100) ⁴⁴	-	-	Effectiveness: ≥21 days after one dose: 33.2% (8.3-51.4); ≥14 days after second dose: 87.9% (78.2-93.2) ⁴⁰	
Sinovac	Chile: 67% (65-69) ²²	-	-	-	Brazil: vaccine effectiveness 1 or 2 doses: 35.1% (-6.6-60.5) ⁴⁵ Chile: 67% (65-69) ²² Brazil: ≥70 years: 41.6% (26.9-53.3); 70-74 years: 61.8% (34.8-77.7); 75-79 years: 48.9% (23.3-66.0); ≥80 years: 28.0% (0.6-47.9) ⁴⁶	-		

* While it is known P.1. and B.1.1.7 were circulating at the time of the study, the extent is unknown based on available surveillance

Vaccine Efficacy/Effectiveness in the Elderly and Against Comorbidities

VACCINE	VACCINE EFFICACY UNLESS OTHERWISE STATED			
	DIABETES	OBESITY	AT RISK FOR SEVERE COVID-19	ELDERLY
AstraZeneca	-	-	76% against symptomatic infection in a sample where 60% had comorbidities, including diabetes, severe obesity or cardiac disease ¹³ (not peer-reviewed)	In ≥65 years: 85% ¹³ (not peer-reviewed) Effectiveness against hospitalisation at 28-34 days after a single dose (pooled analysis of AstraZeneca and Pfizer vaccines) 18-64 years: 85% (68-93); 65-79 years: 79% (17-95); ≥80 years: 81% (65-90) ²⁴ Effectiveness against symptomatic infection in England: ≥70 years: 60% (41-73) ⁴⁷ Effectiveness of single dose against hospitalisation in England: ≥80 years: 73% (60-81) ⁴⁸ Effectiveness in England: Symptomatic infection ≥70 years: 73% (27-90); Hospitalisation ≥80 years: 37% (3-59) ⁴⁹ Hospitalisation following single dose in the UK: ≥80 years: 80.4% (36.4-94.5) ⁵⁰ Single dose in Spain: ≥60 years: 53% (19-72) vs. 18-59 years: 50% (34-62) ²⁶
Gamaleya	-	-	-	Against symptomatic infection in >60 years: 91.8% (67.1-98.3) ¹⁴
Johnson & Johnson	Against moderate to severe/critical: 23.0% (-90.1-69.8) ¹⁵	Against moderate to severe/critical: 65.9% (47.8-78.3) ¹⁵	Against moderate to severe/critical: With any comorbidity: 58.6% (40.6-71.6) ¹⁵ No comorbidity: 68.8% (59.0-76.6) ¹⁵	Against moderate-severe/critical disease ≥28 post vaccination: 18-59 years: 66.1% (53.3-75.8) 60+ years: 66.2% (36.7-83.0) ¹⁵
Moderna	-	-	Against symptomatic infection, comorbidities, including diabetes and obesity: In low risk: 95.1% (89.6-97.7) In high risk: 90.9% (74.7-96.7) ¹⁶	Against symptomatic infection: 18-64 years: 95.6% (90.6-97.9) ≥65 years: 86.4% (61.4-95.2) ¹⁶ Pooled Moderna and Pfizer vaccines against hospitalisation ≥65 years: 94% (49-99) ⁵¹
Pfizer/BioNTech	Effectiveness in Israel: Diabetes or cardiovascular disease: 82% (62-92) ³⁴	-	Against symptomatic infection: With any comorbidity or obesity: 95.3% With no comorbidity: 94.7% ²⁰ Denmark: Infection: 71% (58-80); Hospitalisation: 81% (49-93) ³³	Against symptomatic infection: >55 years: 93.7% (80.6-98.8) >65 years: 94.7% (66.7-99.9); >75 years: 100% (-13.1-100) ²⁰ Effectiveness against hospitalisation 28-34 days after a single dose (pooled analysis of AstraZeneca and Pfizer vaccines): 18-64 years: 85% (68-93); 65-79 years: 79% (17-95); ≥80 years: 81% (65-90) ²⁴ Effectiveness against symptomatic infection in England: ≥70 years: 61% (51-69) ≥80 years: 89% (85-93) ⁴⁷ (not peer reviewed) England 80-83 years: Documented infection: 70.1% (55.1-80.1) Hospital attendance: 78.9% (60.0-89.9); Hospital admission: 75.6% (52.8-87.6) ⁵² Reduction in incidence of infection in vaccinated people aged >60 years and unvaccinated people aged 20-39 years, respectively: Documented infection: 45% versus 28%; Hospitalisation: 68% versus 22% ⁵³ Pooled Moderna and Pfizer vaccines against hospitalisation ≥65 years: 94% (49-99) ⁵¹ Effectiveness in England: Symptomatic infection ≥70 years: 61% (51-69) Hospitalisation ≥80 years: 43% (33-52); Death ≥80 years (vaccine failure vs non-vaccinated): 51% (37-62) ⁴⁹ Effectiveness against hospitalisation in England ≥80 years: Single dose: 81% (76-85) Fully vaccinated: 93% (89-95) ⁴⁸ (not peer reviewed) Effectiveness in Israel: 65-74 years: 82% (63-92); ≥75 years: 82% (61-91) ³⁴ Hospitalisation following single dose in the UK: ≥80 years: 71.4% (43.1-86.2) ⁵⁰ Single dose in Spain: ≥60 years: 76% (55-87) vs. 18-59 years: 85% (74-91) ²⁶ Effectiveness against infection in Denmark: ≥80 years: 77% (50-89) ³³
Sinovac	-	74.9% (53.7-86.4) ²²	Any comorbidity: 48.9% (26.6-64.5) ²²	-
Sinopharm	-	80.7% (56.7-91.4) ²²	-	Effectiveness against symptomatic infection in Bahrain: ≥60 years: 91% (87-94)

Vaccine Efficacy/Effectiveness Against Transmission

There are limitations related to the analysis and comparison of transmission data between studies and vaccines. Criteria for testing vary between studies and may include, for example, random testing, testing at defined intervals, or retrospective serology.

VACCINE	EFFICACY/EFFECTIVENESS AGAINST ASYMPTOMATIC INFECTION	OTHER OUTCOMES
AstraZeneca	<p>Asymptomatic (UK only): 22.2% (-9.9-45.0) Symptomatic and asymptomatic combined (UK, South Africa and Brazil): 54.1% (44.7-61.9)¹² Odds ratio for household contacts of vaccinated health workers vs non-vaccinated health workers testing positive in England: 0.53 (0.43-0.63)⁵⁴</p> <p>Pooled analysis of AstraZeneca and Pfizer/BioNTech in Scotland: Hazard ratio for household contacts of vaccinated health workers vs non-vaccinated health workers testing positive: 0.70 (0.63-0.78)⁵⁵</p> <p>Hazard ratio for single dose in vaccinated vs. unvaccinated care facility residents in England: 0.32 (0.15-0.66)⁵⁶</p>	-
Johnson & Johnson	Asymptomatic: 59.7% (32.8-76.6) ¹⁵	-
Moderna	<p>US: Pooled analysis of Pfizer/BioNTech and Moderna vaccines: 88.7% (68.4-97.1)⁵⁷ Pooled analysis of Pfizer/BioNTech and Moderna vaccines in USA (weekly testing for 13 weeks): 2 weeks after single dose: 80% (59-90); 2 weeks after second dose: 90% (68%-97)⁵⁸</p> <p>Pooled analysis of Pfizer/BioNTech and Moderna vaccines in USA: compared to unvaccinated residents, relative risk of infection in asymptomatic pre-surgical patients >10 days after first dose: 0.21 (0.12-0.37)⁵⁹</p> <p>Following mRNA vaccination in nursing homes in USA, incident cases in <i>unvaccinated</i> residents decreased from 4.3% within 14 days of the first vaccination clinic to 0.3% after 42 days⁶⁰</p>	-
Pfizer/BioNTech	<p>England: 86% (76-97) 7 days after 2 doses; 72% (58-86) 21 days after 1 dose⁶¹ Israel: 75% (72-84) 15-28 days after single dose⁶²; 92% (88-95)³¹ Israel: 91.5% (90.7-92.2)³²</p> <p>USA: Pooled analysis of Pfizer/BioNTech and Moderna vaccines: 88.7% (68.4-97.1)⁵⁷ UK, single dose: 4-fold decrease in risk amongst HCWs ≥12 days post-vaccination⁶³</p> <p>Pooled analysis of Pfizer/BioNTech and Moderna vaccines in US (weekly testing for 13 weeks): 2 weeks after single dose: 80% (59-90); 2 weeks after second dose: 90% (68%-97)⁵⁸</p> <p>Pooled analysis of Pfizer/BioNTech and Moderna vaccines in USA: compared to unvaccinated residents, relative risk of infection in asymptomatic pre-surgical patients >10 days after first dose: 0.21 (0.12-0.37)⁵⁹</p> <p>Following mRNA vaccination in nursing homes in USA, incident cases in <i>unvaccinated</i> residents decreased from 4.3% within 14 days of the first vaccination clinic to 0.3% after 42 days⁶⁰</p> <p>Effectiveness in unvaccinated household contacts of health workers: 2 weeks after first dose: 8.7% (-28.9-35.4); 10 weeks after first dose: 42.9% (22.3-58.1)⁶⁴</p>	<p>Lower viral load in vaccine failure cases 12-37 days after the first dose of vaccine compared to within the first 11 days, indicating potentially lower infectiousness⁶⁵</p> <p>Data from 223 communities in Israel: strong correlation between community vaccination rate and a later decline in infection among children under 16 years of age who were unvaccinated⁶⁶</p> <p>Substantially decreased viral load for infections occurring 12-37 days after the first dose of vaccine in Israel, indicating likely lower infectiousness⁶⁵</p> <p>Detectable transmission in long-term care facilities in Spain reduced by 90% (76-93)⁶⁷</p> <p>Odds ratio for household contacts of vaccinated health workers vs non-vaccinated health workers testing positive in England: 0.51 (0.44-0.59)⁵⁴</p> <p>Pooled analysis of AstraZeneca and Pfizer/BioNTech in Scotland: Hazard ratio for household contacts of vaccinated health workers vs non-vaccinated health workers testing positive: 0.70 (0.63-0.78)⁵⁵</p> <p>Hazard ratio for single dose in vaccinated vs. unvaccinated care facility residents: 0.35 (0.17-0.71)⁵⁶</p>

Mixed Dose Vaccine Safety and Immune Responses

Mixed vaccine schedules (i.e. delivering different types of vaccine for the first and second dose) could be particularly useful to facilitate better protection against variants of concern and enable vaccination programs to continue if a particular vaccine is unavailable.

SCHEDULE	SAFETY	IMMUNE RESPONSES	OTHER OUTCOMES
AstraZeneca followed by Pfizer/BioNTech	Spain: Similar side effects to those receiving 2 doses of the same vaccine; no safety concerns (not peer reviewed) ⁶⁸ UK: Greater systemic side effects (mild-moderate symptoms) following the booster dose than with 2 doses of the same vaccine; no safety concerns ⁶⁹	Spain: ≥8 week dose interval: Stronger immune response following Pfizer/BioNTech than after 2 doses of AstraZeneca vaccine (not peer reviewed) ⁶⁸ UK: 4 week and 12 week dose intervals: Trial underway with immunogenicity data expected in June 2021 ⁶⁹ Germany: 9-12 week dose interval: Significantly stronger immune response following Pfizer/BioNTech than after 2 doses of AstraZeneca and slightly stronger than after 2 doses of Pfizer/BioNTech (not peer reviewed) ⁷⁰	-
Pfizer/BioNTech followed by AstraZeneca	UK: Greater systemic side effects (mild-moderate symptoms) following the booster dose than with 2 doses of the same vaccine; no safety concerns ⁶⁹	UK: 4 week and 12 week dose intervals: Trial underway with immunogenicity data expected in June 2021 ⁶⁹	-
Pfizer/BioNTech or Johnson & Johnson followed by Moderna	-	USA: Trial underway with 12-20 week dose interval ⁷¹	-
AstraZeneca, Moderna and Pfizer/BioNTech	-	Canada: Trial underway mixing and matching all three vaccines with study arms assessing 4 week and 16 week dose intervals ⁷²	-

Adverse Events Following Immunization with WHO EUL Vaccines

Adverse events following immunisation (AEFIs) are any reactions occurring after immunisation. They can be either expected or unexpected. The vaccine may not actually cause the AEFI; it may occur coincidentally as millions of people are being vaccinated so some people may get sick after vaccination but this does not necessarily mean that it is due to the vaccine. Special investigations determine whether they are due to the vaccine. Adverse events of special interest (AESIs) are of scientific and medical concern that are found through active surveillance, that have the potential to be causally associated with a vaccine and that need to be carefully monitored and confirmed by further special studies.

For all injectable vaccines, appropriate medical treatment and supervision should always be readily available in case of an anaphylactic event following administration.

	ASTRAZENECA	MODERNA	PFIZER/BIONTECH	JOHNSON & JOHNSON	SINOPHARM	SINOVAC
Adverse events following immunisation (AEFIs)*	<p>Very common (more than 1 in 10 people): headache, nausea, muscle pain, joint pain, injection site tenderness/ pain/ warmth/ itch, fatigue, malaise, fever, chills</p> <p>Common (between 1 in 10 and 1 in 100 people): injection site swelling/ redness⁷³</p>	<p>Injection site pain (92%)/ swelling (15%)/ redness (10%), fatigue (70%), headache (65%), muscle pain (62%), joint pain (46%), fever (16%), chills (45%), nausea/vomiting (23%), axillary swelling/tenderness (20%)⁷⁴</p>	<p>Very common: headache, muscle pain, joint pain, injection site pain/ swelling, fatigue, fever, chills;</p> <p>Common: nausea, injection site redness⁷³</p> <p>Uncommon (between 1 in 100 and 1 in 1000 people): lymphadenopathy, insomnia, pain in extremity of vaccinated arm, malaise, injection site itch;</p> <p>Rare: (between 1 in 1000 and 1 in 10,000): acute peripheral facial paralysis⁷⁵</p>	<p>Injection site pain/ redness/ swelling, headache, fatigue, muscle pain, nausea, fever⁷⁶</p>	<p>Injection site pain (16%)/ itch (1%)/ swelling (2%)/ redness (1%), fever (4%), fatigue (3%), nausea (1%), headache (1%), diarrhoea (1%), muscle pain (<1%), itch (non-injection site) (1%)⁷⁷</p>	<p>Fatigue (8.3%), fever (3.3%), diarrhoea (0.8%), nausea (1.7%), headache (2.5%), muscle pain (1.7%), injection site pain (10.0%)/ redness (0%)/ swelling (0%)⁷⁸</p>
Adverse events of special interest (AESIs)	<p>Thrombosis with thrombocytopenia syndrome (TTS) (see page 11 for estimated risk); Guillain-Barre syndrome⁷⁹</p>	<p>Myocarditis;⁸⁰ Immune thrombocytopenia (ITP)**⁸¹</p>	<p>USA: Myocarditis or myopericarditis in 7 males aged 14-19 years within 4 days of the second dose – all cases resolved⁸²</p> <p>Israel: Myocarditis estimated to occur in 1 in 3,000 to 1 in 6,000 men aged 16-24 following the second dose – mostly mild and resolved⁸³</p> <p>ITP**⁸¹</p>	<p>TTS (see page 11 for estimated risk)</p>	-	-

*Details for AstraZeneca, Moderna, Pfizer/BioNTech and Johnson & Johnson from product information sheets in SRA countries, based on data from clinical trials; Sinopharm and Sinovac details from published clinical trials

**The ITP cases are mostly without the thrombotic events characteristic of TTS

Serious Adverse Events

Caution is required when comparing safety profiles as definitions and reporting systems vary in trials and in particular phase IV studies

VACCINE	VACCINE SAFETY
AstraZeneca	<p>108 SAEs in 12,282 (0.9%) vaccine recipients and 127 in 11,962 (1.1%) placebo recipients 12 thromboembolic events (4 vaccine; 8 placebo) 7 deaths, all considered unrelated to vaccination (2 vaccine, 5 placebo)¹²</p> <p>US Phase III study: No serious safety concerns involving 32,449 participants¹³ (not peer-reviewed)</p> <p>EMA investigation: possible link between the AstraZeneca vaccine and Thrombosis with Thrombocytopenia Syndrome (TTS) Blood clots affected the brain (central venous sinus thrombosis, CVST) and abdomen (splanchnic vein thrombosis) There have been reports of 169 cases of CVST and 53 cases of splanchnic vein thrombosis in ~34 million vaccinated people in Europe The EMA confirmed the overall benefits of the vaccine in preventing COVID-19 outweigh the risks of side effects⁶</p> <p>Australia: 20 cases of TTS reported in 1,500,000 people vaccinated⁸⁴</p> <p>Several countries have recommended that only older adults should receive the vaccine (including only those aged over 60 years in Germany; over 55 years in France and Canada; over 50 years in Australia; and over 30 years in the UK⁸⁵⁻⁸⁷)</p> <p>EMA has started a review of reports of capillary leak syndrome following 5 cases of this very rare disorder post vaccination⁸⁸</p>
Gamaleya	<p>45 SAEs in 16,427 (0.3%) vaccine recipients and 23 in 5,435 (0.4%) placebo recipients All SAEs were considered unrelated to vaccination 4 deaths, all considered unrelated to vaccination (3 vaccine, 1 placebo)¹⁴</p>
Johnson & Johnson	<p>83 SAEs in 21,895 (0.4%) vaccine recipients and 96 SAEs in 21,888 placebo recipients (0.4%) 19 deaths all considered unrelated to vaccination (3 vaccine, 16 placebo)¹⁵</p> <p>EMA investigation of 8 reports of TTS: possible link between the Johnson & Johnson vaccine and TTS. Most cases occurred in women <60 years of age but specific risk factors have not been confirmed⁷</p> <p>The CDC and FDA have now recommenced the vaccination program in the USA following a thorough safety review⁸⁹</p> <p>15 cases of TTS have been reported in 7.98 million people vaccinated in USA⁵</p>
Moderna	<p>153 SAEs in 15,166 (1.0%) placebo recipients and 147 in 15,185 (1.0%) vaccine recipients 5 deaths considered unrelated to vaccine (2 vaccine, 3 placebo)¹⁶</p> <p>Anaphylaxis reported in the US at a rate of 2.5 per million doses⁹⁰</p> <p>No obvious safety signals among pregnant women who received mRNA COVID-19 vaccines in USA⁹¹</p>
Novavax	<p>SAEs at low levels and similar between vaccine and placebo groups⁹²</p>
Pfizer/BioNTech	<p>SAEs and deaths were low and comparable between vaccine and placebo groups (total 37,586 participants)²⁰ Anaphylaxis reported in the US at a rate of 4.7 per million doses⁹⁰ No obvious safety signals among pregnant women who received mRNA COVID-19 vaccines in USA⁹¹</p>



Risk of Rare Unusual Blood Clotting with Low Blood Platelets (Thrombosis with Thrombocytopenia Syndrome – TTS)

Estimated number of TTS that potentially might occur in Pacific Island Countries if all adults received the AstraZeneca or Johnson & Johnson vaccines, based on most recent official estimate of the adult population in each country and the incidence of these events in Europe and Australia.

COUNTRY	TOTAL POPULATION	ESTIMATED POPULATION AGED 18 YEARS AND OVER*	POTENTIAL NUMBER OF TTS CASES IF ALL ADULTS IN EACH COUNTRY RECEIVED ASTRAZENECA VACCINE**	POTENTIAL NUMBER OF TTS CASES IF ALL ADULTS IN EACH COUNTRY RECEIVED JOHNSON & JOHNSON VACCINE***
American Samoa	55,519	33,311	<1	<1
Cook Islands	15,300	9,180	<1	<1
Federated States of Micronesia	102,300	61,380	0.6-1.1	<1
Fiji	867,000	520,200	5.2-9.6	1.6
French Polynesia	275,918	165,551	1.7-3.1	<1
Guam	159,358	95,615	1.0-1.8	<1
Kiribati	113,400	68,040	0.7-1.3	<1
Marshall Islands	54,900	32,940	<1	<1
Nauru	10,900	6,540	<1	<1
New Caledonia	271,407	162,844	1.6-3.0	<1
Niue	1,611	967	<1	<1
Northern Mariana Islands	53,883	32,330	<1	<1
Palau	18,000	10,800	<1	<1
Papua New Guinea	7,744,700	4,646,820	46.5-86.1	14.6
Samoa	195,979	117,587	1.2-2.2	<1
Solomon Islands	642,000	385,200	3.9-7.1	1.2
Tokelau	1,160	696	<1	<1
Tonga	99,419	59,651	0.6-1.1	<1
Tuvalu	10,507	6,304	<1	<1
Vanuatu	272,173	163,304	1.6-3.0	<1
Wallis and Futuna	11,558	6,935	<1	<1
All Pacific Island Countries	10,976,992	6,586,195	65.9-122.0	20.8

* Based on estimate of 60% of population aged ≥18 years⁹³

** Based on estimates of TTS occurring in ~1 in 100,000 vaccinated adults by the European Medicines Agency and ~1 in 54,000 in Australia^{8,84}

*** Based on estimates of TTS occurring in ~1 in 319,000 vaccinated adults in USA (may be an underestimate as only cerebral venous sinus thrombosis are reported)¹¹

Who Can be Vaccinated Based on WHO SAGE Recommendations?

So far, WHO SAGE have made recommendations for use of AstraZeneca, Moderna, Pfizer/BioNTech, Johnson & Johnson and Sinopharm vaccines:
<https://www.who.int/groups/strategic-advisory-group-of-experts-on-immunization/covid-19-materials>

	ASTRAZENECA	MODERNA	PFIZER/BIONTECH	JOHNSON & JOHNSON	SINOPHARM	SINOVAC
Minimum Age	18 years	18 years	16 years	18 years	18 years	18 years
Maximum Age (SAGE WHO)	None	None	None	None	None	None
Pregnancy	Yes if high priority group & approved by health provider	Yes if high priority group & approved by health provider	Yes if high priority group & approved by health provider	Yes if high priority group & approved by health provider	Yes if high priority group & approved by health provider	Yes if high priority group & approved by health provider
Breastfeeding	Yes	Yes	Yes	Yes	Yes	Yes
Immunocompromised Including HIV	✓	✓	✓	✓	✓	✓
People Previously Infected by SARS-CoV-2 (PCR Confirmed)	Yes, although that person may choose to delay vaccination by 6 months	Yes, although that person may choose to delay vaccination by 6 months	Yes, although that person may choose to delay vaccination by 6 months	Yes, although that person may choose to delay vaccination by 6 months	Yes, although that person may choose to delay vaccination by 6 months	Yes, although that person may choose to delay vaccination by 6 months
History of Anaphylaxis (Severe Allergy)	Yes (unless the allergy is to the vaccine or its components)	Yes (unless the allergy is to the vaccine or its components)	Yes (unless the allergy is to the vaccine or its components)	Yes (unless the allergy is to the vaccine or its components)	Yes (unless the allergy is to the vaccine or its components)	Yes (unless the allergy is to the vaccine or its components)

Vaccine Development Pipeline

WHO has recommended that vaccines adopted by countries have WHO SAGE EUL and/or Stringent Regulatory Approval.

VACCINE TYPE	NUMBER OF VACCINE CANDIDATES AT EACH PHASE OF DEVELOPMENT				
	PRE-CLINICAL	PHASE I/II	PHASE III	PHASE IV	IN USE*
RNA	25	9	2	2	2 (Pfizer/BioNTech, Moderna)
DNA	17	8	2	0	0
Vector (non-replicating)	27	8	3	2	4 (CanSino, Gamaleya, Johnson & Johnson, AstraZeneca)
Vector (replicating)	18	6	0	0	0
Inactivated	8	7	7	2	7 (Sinopharm/BIBP; Sinopharm/WIBP; Sinovac; Bharat; Chumakov; Research Institute for Biological Safety Problems; Shenzhen Kangtai Biological Products)
Live-attenuated	2	1	0	0	0
Protein subunit	75	19	8	1	4 (Vector institute; Anhui Zhifei Longcom Biopharmaceutical Chinese Academy of sciences; Center for Genetic Engineering and Biotechnology; Instituto Finlay de Vacunas Cuba)
Virus-like particle	20	4	1	0	0
Other/unknown	33	5	0	0	0

*Not all vaccines in use have SRA (as recognised by WHO) approval (see Vaccine specifications table and WHO SAGE Emergency Use Listing and prequalification timeline for approval status of vaccines).

Source: London School of Hygiene and Tropical Medicine COVID-19 vaccine tracker.

WHO SAGE Emergency Use Listing and Prequalification Timeline

MANUFACTURER	NAME OF VACCINE	PLATFORM	STATUS OF ASSESSMENT	ANTICIPATED DECISION DATE
Pfizer/BioNTech	BNT162b2/COMIRNATY Tozinameran (INN)	mRNA	Final decision made	Authorised 31/12/20
AstraZeneca	AZD1222	Adenoviral vector	Final decision made	SK Bio: Authorised 15/02/21 EU nodes: Authorised 16/04/21
Serum Institute of India	Covishield (ChAdOx1_nCoV19)	Adenoviral vector	Final decision made	Authorised 15/02/21
Sinopharm/Beijing Institute of Biological Products (BIBP)	SARS-CoV-2 Vaccine (Vero Cell), Inactivated (InCoV)	Inactivated	In progress	Authorised: 07/05/2021
Sinovac	SARS-CoV-2 Vaccine (Vero Cell), Inactivated	Inactivated	In progress	Authorised 01/06/2021
Moderna	mRNA-1273	mRNA	In progress (to use abridged procedure relying on EMA)	Authorised 30/04/2021
Johnson & Johnson	Ad26.COV2.S	Adenoviral vector	Final decision made	Authorised 12/03/21
The Gamaleya National Center	Sputnik V	Adenoviral vector	Clinical and chemistry, manufacturing and control (CMC) review ongoing	Will be determined when all data are submitted
CanSinoBIO	Ad5-nCoV	Adenoviral vector	Rolling data assessment starting May 2021	-
Novavax	NVX-CoV2373	Protein subunit	Expression of interest submitted 23/02/21; Pre-submission meeting to be planned in April	-
CureVac	Zorecimeran	mRNA	Expression of interest submitted 12/04/21	-

Source: WHO Guidance Document: Status of COVID-19 Vaccines within WHO EUL/PQ evaluation process. Available at: <https://www.who.int/teams/regulation-prequalification/eul/covid-19>

Weekly COVID-19 Vaccine Updates
Number 13, 10 June 2021



References

1. Wallace M, Woodworth KR, Gargano JW, et al. The Advisory Committee on Immunization Practices' Interim Recommendation for Use of Pfizer-BioNTech COVID-19 Vaccine in Adolescents Aged 12–15 Years — United States, May 2021. *MMWR Morb Mortal Wkly Rep*. 2021;70(20):749–752. doi:10.15585/mmwr.mm7020e1
2. UK Government Medicines and Healthcare Products Regulatory Agency (MHRA). The MHRA concludes positive safety profile for Pfizer/BioNTech vaccine in 12- to 15-year-olds. 4 June 2021. Available at: <https://www.gov.uk/government/news/the-mhra-concludes-positive-safety-profile-for-pfizerbiontech-vaccine-in-12-to-15-year-olds>.
3. European Medicines Agency (EMA). First COVID-19 vaccine approved for children aged 12 to 15 in EU 28 May 2021. Available at:
4. Government of Canada. Health Canada authorizes use of the Pfizer-BioNTech COVID-19 vaccine in children 12 to 15 years of age - 5 May 2021. Available at:
5. Center for Disease Control (CDC) Advisory Committee on Immunization Practices (ACIP). Thrombosis with thrombocytopenia syndrome (TTS) following Janssen COVID-19 vaccine 23 April 2021. Available at: <https://www.cdc.gov/vaccines/acip/meetings/downloads/slides-2021-04-23/03-COVID-Shimabukuro-508.pdf>.
6. European Medicines Agency (EMA). AstraZeneca's COVID-19 vaccine: EMA finds possible link to very rare cases of unusual blood clots with low blood platelets 7 April 2021. Available at: <https://www.ema.europa.eu/en/news/astrazenecas-covid-19-vaccine-ema-finds-possible-link-very-rare-cases-unusual-blood-clots-low-blood>.
7. European Medicines Agency (EMA). COVID-19 Vaccine Janssen: EMA finds possible link to very rare cases of unusual blood clots with low blood platelets 20 April 2021. Available at: <https://www.ema.europa.eu/en/news/covid-19-vaccine-janssen-ema-finds-possible-link-very-rare-cases-unusual-blood-clots-low-blood>.
8. European Medicines Agency (EMA) press release. AstraZeneca's COVID-19 vaccine: benefits and risks in context 23 April 2021. Available at: <https://www.ema.europa.eu/en/news/astrazenecas-covid-19-vaccine-benefits-risks-context>.
9. Australian Technical Advisory Group on Immunisation (ATAGI). ATAGI update following weekly COVID-19 meeting – 2 June 2021. Available at: <https://www.health.gov.au/news/atagi-update-following-weekly-covid-19-meeting-2-june-2021>.
10. UK Government Medicines and Healthcare Products Regulatory Agency (MHRA). Coronavirus vaccine - weekly summary of Yellow Card reporting 3 June 2021. Available at: <https://www.gov.uk/government/publications/coronavirus-covid-19-vaccine-adverse-reactions/coronavirus-vaccine-summary-of-yellow-card-reporting>.
11. Centers for Disease Control and Prevention (CDC). Selected Adverse Events Reported after COVID-19 Vaccination. Updated 27 May 2021. Available at:
12. Voysey M, Costa Clemens SA, Madhi SA, et al. Single-dose administration and the influence of the timing of the booster dose on immunogenicity and efficacy of ChAdOx1 nCoV-19 (AZD1222) vaccine: a pooled analysis of four randomised trials. *Lancet*. 2021;397(10277):881–891. doi:10.1016/S0140-6736(21)00432-3
13. AstraZeneca press release. AZD1222 US Phase III primary analysis confirms safety and efficacy 25 March 2021. Available at: <https://www.astrazeneca.com/media-centre/press-releases/2021/azd1222-us-phase-iii-primary-analysis-confirms-safety-and-efficacy.html#:~:text=Positive high-level results from,on Monday 22 March 2021.&text=The vaccine was well tolerated, to the vaccine were>.
14. Logunov DY, Dolzhikova I V, Shcheblyakov D V, et al. Safety and efficacy of an rAd26 and rAd5 vector-based heterologous prime-boost COVID-19 vaccine. *Lancet*. 2021.
15. US federal Drug Administration (FDA) Briefing Document Janssen Ad26.CO2V.S Vaccine for the Prevention of COVID-19. Vaccines and Related Biological Products Advisory Committee Meeting February 26, 2021. Available at: <https://www.fda.gov/media/146217/download>.
16. Baden LR, El Sahly HM, Essink B, et al. Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *N Engl J Med*. 2021. doi:10.1056/nejmoa2035389
17. Moderna press release. Moderna Provides Clinical and Supply Updates on COVID-19 Vaccine Program Ahead of 2nd Annual Vaccines Day 13 April 2021. Available at: <https://investors.modernatx.com/news-releases/news-release-details/moderna-provides-clinical-and-supply-updates-covid-19-vaccine>.
18. Moderna press release. Moderna Announces TeenCOVE Study of its COVID-19 Vaccine in Adolescents Meets Primary Endpoint and Plans to Submit Data to Regulators in Early June 25 May 2021. Available at:
19. Heath PT, Galiza EP, Baxter DN, et al. Efficacy of the NVX-CoV2373 Covid-19 Vaccine Against the B.1.1.7 Variant. *medRxiv*. January 2021:2021.05.13.21256639. doi:10.1101/2021.05.13.21256639
20. World Health Organisation (WHO). *Background Document on the mRNA Vaccine BNT162b2 (Pfizer-BioNTech) against COVID-19*; 2021. [https://www.who.int/publications/i/item/background-document-on-mrna-vaccine-bnt162b2-\(pfizer-biontech\)-against-covid-19](https://www.who.int/publications/i/item/background-document-on-mrna-vaccine-bnt162b2-(pfizer-biontech)-against-covid-19).
21. Mahase E. Covid-19: Pfizer reports 100% vaccine efficacy in children aged 12 to 15. *BMJ*. April 2021:n881. doi:10.1136/bmj.n881
22. World Health Organisation Extraordinary meeting of the Strategic Advisory Group of Experts on Immunization (SAGE) – 29 April 2021. Available at: [https://www.who.int/news-room/events/detail/2021/04/29/default-calendar/extraordinary-meeting-of-the-strategic-advisory-group-of-experts-on-immunization-\(sage\)-29-april-2021](https://www.who.int/news-room/events/detail/2021/04/29/default-calendar/extraordinary-meeting-of-the-strategic-advisory-group-of-experts-on-immunization-(sage)-29-april-2021).
23. Palacios R, Batista AP, Albuquerque CSN, et al. Efficacy and Safety of a COVID-19 Inactivated Vaccine in Healthcare Professionals in Brazil: The PROFISCOV Study. *SSRN Electron J*. 2021.
24. Vasileiou E, Simpson CR, Robertson C, et al. Effectiveness of First Dose of COVID-19 Vaccines Against Hospital Admissions in Scotland: National Prospective Cohort Study of 5.4 Million People. *SSRN Electron J*. 2021. doi:10.2139/ssrn.3789264
25. Bernal J, Andrews N, Gower C, et al. Effectiveness of BNT162b2 mRNA vaccine and ChAdOx1 adenovirus vector vaccine on mortality following COVID-19. *Prepr medRxiv*. doi:10.1101/2021.05.14.21257218
26. Martínez-Baz I, Miqueleiz A, Casado I, et al. Effectiveness of COVID-19 vaccines in preventing SARS-CoV-2 infection and hospitalisation, Navarre, Spain, January to April 2021. *Eurosurveillance*. 2021;26(21). doi:10.2807/1560-7917.ES.2021.26.21.2100438
27. University College London press release. Covid-19 vaccine: care home residents gain 62% protection from one dose. 29 March 2021.
28. Pritchard E, Matthews PC, Stoesser N, et al. Impact of vaccination on SARS-CoV-2 cases in the community: a population-based study using the UK's COVID-19 Infection Survey. *medRxiv*. January 2021:2021.04.22.21255913. doi:10.1101/2021.04.22.21255913
29. Daniel W, Nivet M, Warner J, Podolsky DK. Early Evidence of the Effect of SARS-CoV-2 Vaccine at One Medical Center. *N Engl J Med*. March 2021:NEJMc2102153. doi:10.1056/NEJMc2102153
30. Thompson MG, Burgess JL, Naleway AL, et al. Interim Estimates of Vaccine Effectiveness of BNT162b2 and mRNA-1273 COVID-19 Vaccines in Preventing SARS-CoV-2 Infection Among Health Care Personnel, First Responders, and Other Essential and Frontline Workers — Eight U.S. Locations, December 2020–March . *MMWR Morb Mortal Wkly Rep*. 2021;70(13):495–500. doi:10.15585/mmwr.mm7013e3
31. Dagan N, Barda N, Kepten E, et al. BNT162b2 mRNA Covid-19 Vaccine in a Nationwide Mass Vaccination Setting. *N Engl J Med*. February 2021:NEJMoa2101765. doi:10.1056/NEJMoa2101765
32. Haas EJ, Angulo FJ, McLaughlin JM, et al. Impact and effectiveness of mRNA BNT162b2 vaccine against SARS-CoV-2 infections and COVID-19 cases, hospitalisations, and deaths following a nationwide vaccination campaign in Israel: an observational study using national surveillance data. *Lancet*. 2021;397(10287):1819–1829. doi:10.1016/S0140-6736(21)00947-8
33. Emborg H-D, Valentiner-Branth P, Schelde AB, et al. Vaccine effectiveness of the BNT162b2 mRNA COVID-19 vaccine against RT-PCR confirmed SARS-CoV-2 infections, hospitalisations and mortality in prioritised risk groups. *medRxiv*. January 2021:2021.05.27.21257583. doi:10.1101/2021.05.27.21257583
34. Chodick G, Tene L, Rotem RS, et al. The effectiveness of the TWO-DOSE BNT162b2 vaccine: analysis of real-world data. *Clin Infect Dis*. May 2021. doi:10.1093/cid/ciab438
35. Benenson S, Oster Y, Cohen MJ, Nir-Paz R. BNT162b2 mRNA Covid-19 Vaccine Effectiveness among Health Care Workers. *N Engl J Med*. March 2021:NEJMc2101951. doi:10.1056/NEJMc2101951
36. Government of Chile Ministry of Health. Effectiveness of the inactivated CoronaVac vaccine against SARSCoV-2 in Chile Preliminary report April 16 2021.
37. World Health Organisation (WHO). Weekly epidemiological update on COVID-19 - 11 May 2021. Available at: <https://www.who.int/publications/m/item/weekly-epidemiological-update-on-covid-19---11-may-2021>.
38. World Health Organisation (WHO). Tracking SARS-CoV-2 variants. Available at:
39. Emary KRW, Golubchik T, Aley PK, et al. Efficacy of ChAdOx1 nCoV-19 (AZD1222) vaccine against SARS-CoV-2 variant of concern 202012/01 (B.1.1.7): an exploratory analysis of a randomised controlled trial. *Lancet*. March 2021. doi:10.1016/S0140-6736(21)00628-0
40. Bernal J, Andrews N, Gower C, et al. Effectiveness of COVID-19 vaccines against the B.1.617.2 variant. Khub Preprint. Available at: <https://khub.net/documents/135939561/430986542/Effectiveness+of+COVID-19+vaccines+against+the+B.1.617.2+variant.pdf/204c11a4-e02e-112d-db19-b3664107ac42>.
41. Madhi SA, Baillie V, Cutland CL, et al. Safety and efficacy of the ChAdOx1 nCoV-19 (AZD1222) Covid-19 vaccine against the B.1.351 variant in South Africa. *Alex Sigal*. 2021.
42. Shinde V, Bhikha S, Hoosain Z, et al. Efficacy of NVX-CoV2373 Covid-19 Vaccine against the B.1.351 Variant. *N Engl J Med*. May 2021:NEJMoa2103055. doi:10.1056/NEJMoa2103055
43. Kustin T, Harel N, Finkel U, et al. Evidence for increased breakthrough rates of SARS-CoV-2 variants of concern in BNT162b2 mRNA vaccinated individuals. *medRxiv*. January 2021:2021.04.06.21254882. doi:10.1101/2021.04.06.21254882
44. Abu-Raddad LJ, Chemaitelly H, Butt AA. Effectiveness of the BNT162b2 Covid-19 Vaccine against the B.1.1.7 and B.1.351 Variants. *N Engl J Med*. May 2021:NEJMc2104974. doi:10.1056/NEJMc2104974

45. Hitchings MDT, Ranzani OT, Scaramuzzini Torres MS, et al. Effectiveness of CoronaVac in the setting of high SARS-CoV-2 P.1 variant transmission in Brazil: A test-negative case-control study. *medRxiv*. January 2021:2021.04.07.21255081. doi:10.1101/2021.04.07.21255081
46. Ranzani OT, Hitchings M, Nieto MD, et al. Effectiveness of the CoronaVac vaccine in the elderly population during a P.1 variant-associated epidemic of COVID-19 in Brazil: A test-negative case-control study. *medRxiv*. January 2021:2021.05.19.21257472. doi:10.1101/2021.05.19.21257472
47. Bernal JL, Andrews N, Gower C, et al. Early effectiveness of COVID-19 vaccination with BNT162b2 mRNA vaccine and ChAdOx1 adenovirus vector vaccine on symptomatic disease, hospitalisations and mortality in older adults in England. *medRxiv*. January 2021:2021.03.01.21252652. doi:10.1101/2021.03.01.21252652
48. Ismail S, Vilaplana T, Elgohari S, et al. Effectiveness of BNT162b2 mRNA and ChAdOx1 adenovirus vector COVID-19 vaccines on risk of hospitalisation among older adults in England: an observational study using surveillance data. *Khub Prepr* <https://khub.net/documents/135939561/430986542/Effectiveness%20of%20BNT162b2%20mRNA%20and%20ChAdOx1%20adenovirus%20vector%20COVID-19%20vaccines%20on%20risk%20of%20hospitalisation%20among%20older%20adults%20in%20England.pdf/9e18c525-dde6-5ee4>.
49. Lopez Bernal J, Andrews N, Gower C, et al. Effectiveness of the Pfizer-BioNTech and Oxford-AstraZeneca vaccines on covid-19 related symptoms, hospital admissions, and mortality in older adults in England: test negative case-control study. *BMJ*. May 2021:n1088. doi:10.1136/bmj.n1088
50. Hyams C, Marlow R, Maseko Z, et al. Assessing the Effectiveness of BNT162b2 and ChAdOx1nCoV-19 COVID-19 Vaccination in Prevention of Hospitalisations in Elderly and Frail Adults: A Single Centre Test Negative Case-Control Study. *SSRN Electron J*. 2021. doi:10.2139/ssrn.3796835
51. Tenforde MW, Olson SM, Self WH, et al. Effectiveness of Pfizer-BioNTech and Moderna Vaccines Against COVID-19 Among Hospitalized Adults Aged ≥65 Years — United States, January–March 2021. *MMWR Morb Mortal Wkly Rep*. 2021;70(18). doi:10.15585/mmwr.mm7018e1
52. Mason T, Whitston M, Hodgson J, et al. Effects of BNT162b2 mRNA vaccine on Covid-19 infection and hospitalisation among older people: matched case control study for England. *medRxiv*. January 2021:2021.04.19.21255641. doi:10.1101/2021.04.19.21255641
53. Rossman H, Shilo S, Meir T, Gorfine M, Shalit U, Segal E. COVID-19 dynamics after a national immunization program in Israel. *Nat Med*. April 2021. doi:10.1038/s41591-021-01337-2
54. Harris R, Hall J, Zaidi A, Andrews N, Dunbar J, Dabrera G. Impact of vaccination on household transmission of SARS-CoV-2 in England. *Preprint*. <https://khub.net/documents/135939561/390853656/Impact+of+vaccination+on+household+transmission+of+SARS-COV-2+in+England.pdf/35bf4bb1-6ade-d3eb-a39e-9c9b25a8122a?i=1619601878136>.
55. V Shah AS, Gribben C, Bishop J, et al. Effect of vaccination on transmission of COVID-19: an observational study in healthcare workers and their households. *medRxiv*. January 2021:2021.03.11.21253275. doi:10.1101/2021.03.11.21253275
56. Shrotri M, Krutikov M, Palmer T, et al. Vaccine effectiveness of the first dose of ChAdOx1 nCoV-19 and BNT162b2 against SARS-CoV-2 infection in residents of Long-Term Care Facilities (VIVALDI study). *medRxiv*. January 2021:2021.03.26.21254391. doi:10.1101/2021.03.26.21254391
57. Pawlowski C, Lenehan P, Puranik A, et al. FDA-authorized COVID-19 vaccines are effective per real-world evidence synthesized across a multi-state health system. *medRxiv*. January 2021:2021.02.15.21251623. doi:10.1101/2021.02.15.21251623
58. Thompson MG, Burgess JL, Naleway AL, et al. Interim Estimates of Vaccine Effectiveness of BNT162b2 and mRNA-1273 COVID-19 Vaccines in Preventing SARS-CoV-2 Infection Among Health Care Personnel, First Responders, and Other Essential and Frontline Workers
59. Tande AJ, Pollock BD, Shah ND, et al. Impact of the COVID-19 Vaccine on Asymptomatic Infection Among Patients Undergoing Pre-Procedural COVID-19 Molecular Screening. *Clin Infect Dis*. March 2021. doi:10.1093/cid/ciab229
60. White EM, Yang X, Blackman C, Feifer RA, Gravenstein S, Mor V. Incident SARS-CoV-2 Infection among mRNA-Vaccinated and Unvaccinated Nursing Home Residents. *N Engl J Med*. May 2021:NEJMc2104849. doi:10.1056/NEJMc2104849
61. Hall VJ, Foulkes S, Saei A, et al. Effectiveness of BNT162b2 mRNA Vaccine Against Infection and COVID-19 Vaccine Coverage in Healthcare Workers in England, Multicentre Prospective Cohort Study (the SIREN Study). *SSRN Electron J*. 2021. doi:10.2139/ssrn.3790399
62. Amit S, Regev-Yochay G, Afek A, Kreiss Y, Leshem E. Early rate reductions of SARS-CoV-2 infection and COVID-19 in BNT162b2 vaccine recipients. *Lancet*. 2021;397(10277):875-877. doi:10.1016/S0140-6736(21)00448-7
63. Michael Weekes, Nick K Jones, Lucy Rivett, et al. Single-dose BNT162b2 vaccine protects against asymptomatic SARS-CoV-2 infection. *Authoria*. February 24, 2021. doi:10.22541/au.161420511
64. Salo J, Hägg M, Kortelainen M, et al. The indirect effect of mRNA-based Covid-19 vaccination on unvaccinated household members. *medRxiv*. January 2021:2021.05.27.21257896. doi:10.1101/2021.05.27.21257896
65. Levine-Tiefenbrun M, Yelin I, Katz R, et al. Initial report of decreased SARS-CoV-2 viral load after inoculation with the BNT162b2 vaccine. *Nat Med*. 2021;27(5):790-792. doi:10.1038/s41591-021-01316-7
66. Milman O, Yelin I, Aharoni N, et al. SARS-CoV-2 infection risk among unvaccinated is negatively associated with community-level vaccination rates. *medRxiv*. January 2021:2021.03.26.21254394. doi:10.1101/2021.03.26.21254394
67. De Salazar PM, Link N, Lamarca K, Santillana M. High coverage COVID-19 mRNA vaccination rapidly controls SARS-CoV-2 transmission in Long-Term Care Facilities. *medRxiv*. January 2021:2021.04.08.21255108. doi:10.1101/2021.04.08.21255108
68. Government of Spain Ministry of Health and Innovation. The combined use of AstraZeneca and Pfizer vaccines against SARS-CoV-2 offers a powerful immune response. Press release 18 May 2021. Available at: <https://www.isciii.es/Noticias/Noticias/Paginas/Noticias/Presentación-resultados-preliminares-Combivac.aspx>.
69. Shaw RH, Stuart A, Greenland M, Liu X, Van-Tam JSN, Snape MD. Heterologous prime-boost COVID-19 vaccination: initial reactivity data. *Lancet*. 2021;397(10289):2043-2046. doi:10.1016/S0140-6736(21)01115-6
70. Saariand University press release. Study: Particularly clear immune response with the vaccine combination of Astrazeneca and Biontech - 6 June 2021. Available at: <https://www.uni-saarland.de/en/university/news/news-article/nr/23581.html>.
71. United States National Institutes of Health (NIH) press release. NIH clinical trial evaluating mixed COVID-19 vaccine schedules begins. Available at: <https://clinicaltrials.gov/ct2/show/NCT04894435?term=mixed+schedule%2C+covid19%2C+vaccine&draw=2&rank=1>.
72. US National Library of Medicine ClinicalTrials.gov. Mix and Match of the Second COVID-19 Vaccine Dose for Safety and Immunogenicity (MOSAIC). Available at: <https://clinicaltrials.gov/ct2/show/NCT04894435?term=mixed+schedule%2C+covid19%2C+vaccine&draw=2&rank=1>.
73. Australian Government Therapeutic Goods Administration (TGA). Australian Product Information - COVID-19 Vaccine AstraZeneca (ChAdOx1-S) solution for injection. Available at: <https://www.ebs.tga.gov.au/ebs/picmi/picmirepository.nsf/pdf?OpenAgent&id=CP-2021-PI-01194-1&d=202105261016933>.
74. US Federal Drug Administration (FDA). Fact sheet for healthcare providers administering vaccine (vaccination providers) emergency use authorisation (EUA) of the Moderna COVID-19 vaccine to prevent coronavirus disease 2019 (COVID-19). Available at: <https://www.modernatx.com/covid19vaccine-eua/eua-fact-sheet-providers.pdf>.
75. Australian Government Therapeutic Goods Administration (TGA). Australian Product Information - Comirnaty (BNT162b2 [mRNA]) COVID-19 Vaccine. Available at: <https://www.ebs.tga.gov.au/ebs/picmi/picmirepository.nsf/pdf?OpenAgent&id=CP-2021-PI-01092-1>.
76. US Federal Drug Administration (FDA). Fact sheet for healthcare providers administering vaccine (vaccination providers). Emergency use authorisation (EUA) of the Janssen COVID-19 vaccine to prevent coronavirus disease 2019 (COVID-19). Available at: <https://www.janssenlabels.com/emergency-use-authorization/Janssen+COVID-19+Vaccine-HCP-fact-sheet.pdf>.
77. Xia S, Zhang Y, Wang Y, et al. Safety and immunogenicity of an inactivated SARS-CoV-2 vaccine, BBIBP-CoVr: a randomised, double-blind, placebo-controlled, phase 1/2 trial. *Lancet Infect Dis*. 2021;21(1):39-51. doi:10.1016/S1473-3099(20)30831-8
78. Zhang Y, Zeng G, Pan H, et al. Safety, tolerability, and immunogenicity of an inactivated SARS-CoV-2 vaccine in healthy adults aged 18–59 years: a randomised, double-blind, placebo-controlled, phase 1/2 clinical trial. *Lancet Infect Dis*. 2021;21(2):181-192. doi:10.1016/S1473-3099(20)30843-4
79. European Medicines Agency (EMA). Meeting highlights from the Pharmacovigilance Risk Assessment Committee (PRAC) 3-6 May 2021. Available at: <https://www.ema.europa.eu/en/news/meeting-highlights-pharmacovigilance-risk-assessment-committee-prac-3-6-may-2021>.
80. World Health Organisation (WHO). COVID-19 subcommittee of the WHO Global Advisory Committee on Vaccine Safety (GACVS) reviews cases of mild myocarditis reported with COVID-19 mRNA vaccines. Available at: <https://www.who.int/news/item/26-05-2021-gacvs-mycarditis-reported-with-covid-19-mrna-vaccines>.
81. Welsh KJ, Baumbblatt J, Chege W, Goud R, Nair N. Thrombocytopenia including immune thrombocytopenia after receipt of mRNA COVID-19 vaccines reported to the Vaccine Adverse Event Reporting System (VAERS). *Vaccine*. 2021;96(5):534-537. doi:10.1016/j.vaccine.2021.04.054
82. Marshall M, Ferguson ID, Lewis P, et al. Symptomatic Acute Myocarditis in Seven Adolescents Following Pfizer-BioNTech COVID-19 Vaccination. *Pediatrics*. June 2021:e2021052478. doi:10.1542/peds.2021-052478
83. Vogel G. Israel reports link between rare cases of heart inflammation and COVID-19 vaccination in young men. *Science (80-)*. June 2021. doi:10.1126/science.abj7796
84. Australian Technical Advisory Group on Immunisation (ATAGI). ATAGI update following weekly COVID-19 meeting – 19 May 2021. Available at: <https://www.health.gov.au/news/atagi-update-following-weekly-covid-19-meeting-19-may-2021>.
85. Government of Canada National Advisory Committee on Immunization (NACI) statement. NACI rapid response: Recommended use of AstraZeneca COVID-19 vaccine in younger adults.

86. UK Government Department of Health and Social Care. Joint Committee on Vaccination and Immunisation (JCVI) statement on use of the AstraZeneca COVID-19 vaccine: 7 April 2021.
87. Australian Technical Advisory Group on Immunisation (ATAGI). ATAGI reinforce recommendations on use of COVID-19 vaccines following review of vaccine safety data and benefits 23 April 2021. Available at: <https://www.health.gov.au/news/atagi-reinforce-recommendations-on-use-of-covid-19-vaccines-following-review-of-vaccine-safety-data-and-benefits>.
88. European Medicines Agency (EMA). Meeting highlights from the Pharmacovigilance Risk Assessment Committee (PRAC) 6-9 April 2021. Available at: <https://www.ema.europa.eu/en/news/meeting-highlights-pharmacovigilance-risk-assessment-committee-prac-6-9-april-2021>.
89. US Food and Drug Administration (FDA) press release. FDA and CDC Lift Recommended Pause on Johnson & Johnson (Janssen) COVID-19 Vaccine Use Following Thorough Safety Review 23 April 2021. Available at: <https://www.fda.gov/news-events/press-announcements/fda-and-cdc-lift-recommended-pause-johnson-johnson-janssen-covid-19-vaccine-use-following-thorough>.
90. Shimabukuro TT, Cole M, Su JR. Reports of Anaphylaxis After Receipt of mRNA COVID-19 Vaccines in the US—December 14, 2020–January 18, 2021. *JAMA*. 2021;325(11):1101. doi:10.1001/jama.2021.1967
91. Shimabukuro TT, Kim SY, Myers TR, et al. Preliminary Findings of mRNA Covid-19 Vaccine Safety in Pregnant Persons. *N Engl J Med*. April 2021:NEJMoa2104983. doi:10.1056/NEJMoa2104983
92. Novavax COVID-19 Vaccine Demonstrates 89.3% Efficacy in UK Phase 3 Trial. Novavax press release 28 January 2021.
93. United Nations Children's Fund (UNICEF), Pacific Office. Situation Analysis of Children in the Pacific Island Countries. December 2017.

Acknowledgements

This document was compiled in collaboration with the WHO South Pacific Division of Pacific Technical Support (DPS), Suva, Fiji

Compiled by Dr John Hart

Technical leads: Professor Fiona Russell and Professor Kim Mulholland

Quality checks: Professor Julie Bines, Associate Professor Nigel Crawford and Associate Professor Margie Danchin

Other resources on COVID-19 vaccines:

WHO COVID-19 vaccines website: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/covid-19-vaccines>

EMA COVID-19 vaccines website: <https://www.ema.europa.eu/en/human-regulatory/overview/public-health-threats/coronavirus-disease-covid-19/treatments-vaccines/covid-19-vaccines>

To subscribe to receive the weekly updates, please email kase.anderson@unimelb.edu.au

Murdoch Children's Research Institute
50 Flemington Rd, Parkville
Victoria 3052 Australia
ABN 21 006 566 972

Weekly COVID-19 Vaccine Updates
Number 13, 10 June 2021

